

Identities in Representations of the Hecke Algebra

S. Loesch

*C.E.A.-SACLAY, Service de Physique Théorique,
F-91191 Gif sur Yvette Cedex, France*

Y.-K. Zhou

*School of Mathematical Sciences, ANU,
Canberra ACT 0200, Australia*

and

J.-B. Zuber

*C.E.A.-SACLAY, Service de Physique Théorique ,
F-91191 Gif sur Yvette Cedex, France*

New identities on traces of representations of the Hecke algebra on the spaces of paths on graphs are presented. These identities are relevant in the computation of partition functions with fixed boundary conditions and of two-point correlation functions in a class of lattice models. They might also help to determine the Boltzmann weights of these (allegedly) integrable models.

1. Introduction

This paper is devoted to the study of identities satisfied by the Boltzmann weights of some two-dimensional integrable lattice models. These models are originally built from certain representations of the $sl(k)$ algebras, or rather from their quantum deformations. Imposing that their (‘face’) transfer matrices commute with the action of the quantum group $\mathcal{U}_q sl(k)$ implies that these transfer matrices are generated by some representation of the Hecke algebra (in fact a quotient of the Hecke algebra by a condition depending on k , see below) [1]. Thus the original problem of building integrable models of that type is rephrased as the algebraic study of representations of the Hecke algebra.

In the graph models considered here, that are known to have a critical regime described by minimal W_k conformal theories, these representations are supported by the space of paths on a graph [2]. Which graph is adequate for this construction remains to be fully understood, but one may start from the well-known ‘basic’ case of the weight lattice of $sl(k)$ truncated at some level. A few other graphs have been discovered in previous investigations [3–6], more have been conjectured lately [7], but the determination of their Boltzmann weights and the verification that they satisfy the relations of the Hecke algebra is a painful, case-by-case analysis [6]. It has been observed recently, however, that these weights seem to satisfy general identities: traces of products of these weights take a simple, universal (i.e. graph independent) expression. The aim of the present work was thus to understand the origin and the form of these identities and possibly to exploit them for a general determination of the Boltzmann weights.

The present work is also connected with considerations on conformal field theories. On the one hand, the traces that we are considering yield particular contributions to the partition functions of the lattice models with fixed boundary conditions. Along the line explored by previous authors [8,5], these partition functions are known to enjoy algebraic properties that match those of the corresponding conformal field theories [9]. On the other hand, these traces may also be regarded as particular contributions to two-point functions of certain lattice operators. The universality properties of correlation functions with respect to the choice of representation of the Hecke algebra have also consequences on properties of structure constants of the operator algebra of the corresponding conformal field theories: see [10] for a discussion in the case of $sl(2)$.

In sect. 2 we recall the context and notations following previous references. We also state the main results and conjectures of this paper. Section 3 presents the elements of proof that we have collected.

2. The Hecke Algebra. Identities on traces.

2.1. Notations on the Hecke algebra

The Hecke algebra $\mathcal{H}_L(q)$ is a q -deformation of the algebra of the symmetric group S_L . It is generated by the identity 1 and by $g_i, i = 1, \dots, L-1$, the q -analogues of the transpositions $(i, i+1)$ generating the symmetric group S_L , that satisfy

$$\begin{aligned} g_i^2 &= 1 + (q - q^{-1})g_i \\ g_i g_j &= g_j g_i \quad |i - j| \geq 2; \\ g_i g_{i+1} g_i &= g_{i+1} g_i g_{i+1} . \end{aligned} \tag{2.1}$$

Alternatively $\mathcal{H}_L(q)$ is generated by the generators $U_i = q - g_i$ satisfying

$$\begin{aligned} U_i^2 &= (q + q^{-1})U_i \\ U_i U_j &= U_j U_i \quad |i - j| \geq 2; \\ U_i U_{i+1} U_i - U_i &= U_{i+1} U_i U_{i+1} - U_{i+1} \end{aligned} \tag{2.2}$$

We are interested in representations of a quotient of the Hecke algebra by an additional condition. This comes about in the search of lattice integrable models whose diagonal-to-diagonal transfer matrix commutes with the action of the quantum group $\mathcal{U}_q sl(k)$. The commutant of that quantum group in the representation space $[k]^{\otimes L}$ of the L -th power of the k -dimensional fundamental representation is generated by a certain k -dependent quotient of the Hecke algebra $\mathcal{H}_L(q)$.

The extra condition relevant to describe the commutant of $\mathcal{U}_q sl(k)$ is easier to express in terms of the g generators. For any permutation σ of S_{k+1} , write a minimal decomposition into transpositions of adjacent elements: $\sigma = \prod_{i \in I_\sigma} \sigma_i$ and denote by g_σ the corresponding product of generators of the Hecke algebra $g_\sigma = \prod_{i \in I_\sigma} g_i$. Then the condition reads [1]

$$\sum_{\sigma \in S_{k+1}} (-q)^{-|I_\sigma|} g_\sigma = 0 . \tag{2.3}$$

See also [11] for a discussion of this condition.

2.2. Representations on path spaces of graphs

We consider representations of this (quotient of the) Hecke algebra on the space of paths of graphs.

When dealing with $\mathcal{U}_q sl(k)$ the basic example of graph is given by the system of integrable weights of the affine algebra $\widehat{sl(k)}_{n-k}$ at a certain level $n-k$, if $q = \exp \frac{i\pi}{n}$. This system of weights is the truncated weight lattice shifted by the vector $\mathbf{1} := \sum_1^{k-1} \Lambda_\ell$

$$\mathcal{P}_{++}^{(n)} = \{\lambda = \lambda_1 \Lambda_1 + \cdots + \lambda_{k-1} \Lambda_{k-1} | \lambda_\alpha \in \mathbb{N}, \lambda_\alpha \geq 1, \lambda_1 + \cdots + \lambda_{k-1} \leq n-1\} . \quad (2.4)$$

We also need the k (linearly dependent) vectors e_i

$$\begin{aligned} e_1 &= \Lambda_1, & e_\alpha &= \Lambda_\alpha - \Lambda_{\alpha-1}, & i &= 2, \dots, k-1, & e_k &= -\Lambda_{k-1} \\ e_1 + e_2 + \cdots + e_k &= 0, \end{aligned} \quad (2.5)$$

endowed with the scalar product

$$(e_\alpha, e_\beta) = \delta_{\alpha\beta} - \frac{1}{k} \quad \alpha, \beta = 1, \dots, k . \quad (2.6)$$

We then consider the fusion coefficients $N_{\lambda\mu}^\nu$ [12] and the associative, commutative algebra that has these coefficients as structure constants. The set of matrices N_λ , of non negative integer entries $(N_\lambda)_\mu^\nu = N_{\lambda\mu}^\nu$, for $\lambda, \mu, \nu \in \mathcal{P}_{++}^{(n)}$, forms the regular representation of this fusion algebra

$$N_\lambda N_\mu = N_{\lambda\mu}^\nu N_\nu . \quad (2.7)$$

If we denote for short $G_\ell = N_{\mathbf{1} + \Lambda_\ell}$ the fusion matrix for the ℓ -th fundamental representation of weight Λ_ℓ (or rather $\mathbf{1} + \Lambda_\ell$ in the notation of (2.4)), it is known that the set $\{N_\lambda\}$ is polynomially generated by the G 's

$$N_\lambda = \mathcal{P}_\lambda(G_1, G_2, \dots, G_{k-1}) . \quad (2.8)$$

The matrices G_ℓ , $\ell = 1, \dots, k-1$ may be regarded as the adjacency matrices of a collection of graphs $\mathcal{G}_\ell$. Only $\mathcal{G}_1$ will be relevant for the considerations of the present paper. In that basic case we shall refer to it as $\mathcal{A}^{(n)}$: it is defined by its vertices, the weights of (2.4), and its edges between neighbouring vertices on the weight lattice oriented along the vectors e_i .

Beside this “basic” case, the other graphs $\mathcal{G}_\ell$ that are known may be recast in that same framework by noticing that their matrices $G_1, G_2, \dots, G_{k-1}$ generate by the same

polynomials as in (2.8) another integrally valued representation $N_\lambda^{(\mathcal{G})}$ of the fusion algebra (2.7) (see [4], where these matrices $(N_\lambda^{(\mathcal{G})})_{ab}$ were denoted $(V_\lambda)_{ab}$.)

In the following, we shall also extend the notation G_ℓ to $G_0 = G_k = \mathbb{I}$, (the identity matrix), $G_\ell = 0$ if $\ell > k$.

One then considers the space $\mathcal{P}^{(L)}$ of paths of length L on the graph $\mathcal{G}_1$, i.e. the space spanned by the states $\{|a_0 \cdots a_L\rangle\}$

$$|a_0 \cdots a_L\rangle = (G_1)_{a_0 a_1} (G_1)_{a_1 a_2} \cdots (G_1)_{a_{L-1} a_L} |a_0\rangle \otimes |a_1\rangle \otimes \cdots |a_L\rangle, \quad (2.9)$$

where $\langle a|b\rangle = \delta_{ab}$ and the matrix elements of the adjacency matrix G_1 of the graph $\mathcal{G}_1$ ensure that consecutive vertices a_i, a_{i+1} along the path are adjacent on the graph. (In this paper, for the sake of simplicity, we consider only graphs $\mathcal{G}_1$ with no double edge: $(G_1)_{ab} = 0$ or 1 .)

When the graph under consideration is one of the basic ones introduced above, it has been shown that the space of paths $\mathcal{P}^{(L)}$ supports a representation of the quotient of the Hecke algebra $\mathcal{H}_L(q)$ given by the following formula [13]

$$U_i |a_0 a_1 \cdots a_i \cdots a_L\rangle = \sum_{a'_i} |a_0 a_1 \cdots a'_i \cdots a_L\rangle \begin{array}{c} a_{i+1} \\ \diamond \\ a_{i-1} \end{array} a_i, \quad i = 1, \dots, L-1,$$

with $\begin{array}{c} a_{i+1} \\ \diamond \\ a_{i-1} \end{array} a_i = (1 - \delta_{\alpha\beta}) \frac{[s_{\alpha\beta}(a_i) s_{\alpha\beta}(a'_i)]^{\frac{1}{2}}}{s_{\alpha\beta}(a_{i-1})}.$ (2.10)

where $a_i = a_{i-1} + e_\alpha$, $a_{i+1} = a_i + e_\beta$ and the function $s_{\alpha\beta}$ is defined as $s_{\alpha\beta}(b) = \sin[(\pi/n)(e_\alpha - e_\beta, b)]$. Here and in the following, the operators U are regarded as acting on vertical chains $a_0 \cdots a_L$ and *from right to left*, in agreement with the usual convention on composition of operators.

For other graphs, subject to additional conditions that are not yet fully clarified, the space of paths also supports a representation of the Hecke algebra. Only in the case of $\widehat{sl(2)}$ in which the relevant graphs turn out to be *ADE* Dynkin diagrams, the general expression of the U 's has been found in terms of the Perron-Frobenius eigenvector $\psi^{(1)}$ of G_1 [2]

$$\begin{array}{c} c \\ \diamond \\ a \end{array} b' = \delta_{ac} \frac{(\psi_b^{(1)} \psi_{b'}^{(1)})^{\frac{1}{2}}}{\psi_a^{(1)}}. \quad (2.11)$$

2.3. Traces

Given a representation of the Hecke algebra of the previous type, namely on paths on a graph $\mathcal{G}_1$, we need the trace of an arbitrary element x of that algebra. As the elements of the algebra do not affect the end points of the path, we may in fact compute the traces in the subspaces of paths with fixed ends a_0 and a_L . The trace tr of the element x of the Hecke algebra $\mathcal{H}_L$ with fixed L is thus a matrix defined as ¹

$$(\text{tr}(x))_{a_0 a_L} := \sum_{a_1, a_2, \dots, a_{L-1}} \langle a_0 a_1 \cdots a_{L-1} a_L | x | a_0 a_1 \cdots a_{L-1} a_L \rangle . \quad (2.12)$$

In words, we are computing the trace of the element x in the subspace $\mathcal{P}_{a_0 a_L}^{(L)}$ of paths of length L going from height a_0 to height a_L . The most important property of the trace (2.12) that we shall need below is its cyclicity : $\text{tr}(xy) = \text{tr}(yx)$.

We are mainly interested in computing expressions of the form

$$\begin{aligned} (Z_L) &= \text{tr} U_1 U_2 \cdots U_{L-1} \\ \text{or} \quad (\tilde{Z}_L) &= \text{tr} g_1 g_2 \cdots g_{L-1} , \end{aligned} \quad (2.13)$$

which may be represented pictorially as in Fig. 1, with each diamond standing for a U or a g , depending on the case.




Fig. 1: Pictorial representation of (2.13). A summation over $a_1, \dots, a_{L-1}$ is understood

¹ The normalization constant is sometimes fixed to be $1/(\gamma_1^{(1)})^L$ so as to make the “Markov trace” $\text{tr}_M(x) := \sum_{a_0, a_L} \psi_{a_0}^{(1)*} \psi_{a_L}^{(1)} (\text{tr}(x))_{a_0 a_L}$ independent of $L > L'$, if x belongs to the algebra generated by $U_1, \dots, U_{L'-1}$. Here and in the following, $\psi^{(1)}$ stands for the Perron-Frobenius eigenvector common to all the G matrices. We shall not make this choice in this paper.

We recall that the local face transfer matrix at site i of the integrable model built out of this representation of the Hecke algebra is a linear combination of U_i (or g_i) and the identity operator [2]. Thus expressions (2.13) are particular cases of what has to be computed in the evaluation of the partition function of the model.

If $L = 1$, the element x is necessarily the identity, the summation in (2.12) is void and we set $Z_1 = \tilde{Z}_1 = G_1$ in agreement with (2.9). Also it is easily verified that the computation of traces where some of the U 's or g 's are skipped reduces to the matrix product of lower ones times possibly G 's; for example

$$\begin{aligned} (\text{tr } U_1 U_3)_{a_0 a_4} &= \sum_{a_1, \dots, a_3, a'_1, \dots, a'_3} \langle a_0 \dots a_4 | U_1 | a_0 a'_1 \dots a'_3 a_4 \rangle \langle a_0 a'_1 \dots a'_3 a_4 | U_3 | a_0 \dots a_4 \rangle \\ &= \sum_{a_1, a_2, a_3} \langle a_0 a_1 a_2 | U_1 | a_0 a_1 a_2 \rangle \langle a_2 a_3 a_4 | U_3 | a_2 a_3 a_4 \rangle = (Z_2 \cdot Z_2)_{a_0 a_4} \end{aligned} \quad (2.14)$$

since U_i “sees” only heights a_{i-1} , a_i and a_{i+1} and affects only a_i . Likewise $\text{tr } U_1 U_4 = Z_2 G_1 Z_2$. A more complicated example is given by (Fig.2).

$$\text{tr } (U_2 U_4 U_5 U_8 U_{11}) = Z_1 Z_2 Z_3 Z_1 Z_2 Z_1^2 Z_2 . \quad (2.15)$$

Notice that the labelling by L of Z_L , $\tilde{Z}_L$ implies that we are dealing with paths of length L from a_0 to a_L , and the expression (2.15) has indeed the grading 13, if each Z_ℓ is counted for ℓ .




Fig. 2: Pictorial representation of (2.15). The squares in solid line indicate the action of a U , those in dotted line stand for the action of 1.

2.4. Properties of the Z and $\tilde{Z}$

We show now that Z and $\tilde{Z}$ are matrices that commute with the matrices G_ℓ and may be expressed in terms of them. Moreover, we establish for low values of k and L , and conjecture in general, some recursion formulae on L for these expressions.

a) First we claim that the Z and $\tilde{Z}$ matrices satisfy

$$Z_L^{(\mathcal{G})} = \sum_{\lambda \in \mathcal{P}_{++}^{(n)}} z_L^{(\lambda)} N_\lambda^{(\mathcal{G})} \quad (2.16)$$

with the same matrices $N_\lambda^{(\mathcal{G})}$ as introduced above and a *universal* coefficient $z_L^{(\lambda)}$, and a similar relation involving coefficients $\tilde{z}_L^{(\lambda)}$ for $\tilde{Z}_L$. Here as above, ‘universal’ means independent of the graph $\mathcal{G}_1$ and depending only on k, n, L ; we shall see below that the dependence of $z_L^{(\lambda)}$ on k, n is just embodied in the restriction of the summation over λ .

Consider first the case of the “basic” graphs $\mathcal{A}^{(n)}$. Then the matrix $N_\lambda^{(\mathcal{A})}$ is nothing else than the fusion matrix N_λ . As the proof of (2.16) is a bit lengthy and technical, it has been relegated to an appendix.

As mentionned in the Introduction, the fact that the traces (2.13) may be reduced through fusion matches the same property of partition functions with fixed boundary conditions in conformal field theories [9]. Note that relation (2.16) allows to determine the coefficient $z_L^{(\lambda)}$ as

$$z_L^{(\lambda)} = (Z_L^{(\mathcal{A})})_{\mathbf{1}\lambda} , \quad (2.17)$$

for the basic graph, and a similar relation for $\tilde{Z}_L^{(\lambda)}$. This is because $N_{\lambda\mathbf{1}}^\mu = \delta_{\lambda\mu}$. Thus (2.16) and (2.17) imply

$$(Z_L^{(\mathcal{A})})_{\mu\nu} = (Z_L^{(\mathcal{A})})_{\mathbf{1}\lambda} N_{\lambda\mu}^\nu , \quad (2.18)$$

The algebraic interpretation of (2.18) is that the only irreducible representations of the Hecke algebra are those on paths on the basic graph running from $\mathbf{1}$ to some weight λ and that the fusion coefficients $N_{\mu\nu}^\lambda$ yield the multiplicities in the decomposition of the more general representation on paths from μ to ν [14,9].

For other graphs $\mathcal{G}_1$, we have no direct proof of the general validity of (2.16), but convergent evidences. First it can be proved for $k = 2$, using the explicit Boltzmann weights (2.11). Also as we shall see below, for $k = 3$, very explicit forms will be given to the Z_L based on the only defining relations of the Hecke algebra, thus valid for all graphs, that agree with (2.16). Finally the same interpretation may be given to the integer entries $\left(N_\lambda^{(\mathcal{G})}\right)_{ab}$ of the matrices $N_\lambda^{(\mathcal{G})}$ as multiplicities in the decomposition of the representation carried by the paths on $\mathcal{G}_1$ running between a and b

$$\left(Z_L^{(\mathcal{G})}\right)_{ab} = \left(Z_L^{(\mathcal{A})}\right)_{\mathbf{1}\lambda} \left(N_\lambda^{(\mathcal{G})}\right)_{ab} . \quad (2.19)$$

b) Eq. (2.8) and (2.16) imply that Z_L and $\tilde{Z}_L$ may be expressed in terms of the fundamental matrices G_ℓ , $\ell = 1, \dots, k-1$. In fact we have formulae that enable us to determine these expressions recursively.

For this we make use of Chebychev polynomials of second type

$$P_\ell(\beta) = \frac{\sin(\ell+1)\theta}{\sin \theta} \quad \beta = 2 \cos \theta. \quad (2.20)$$

(We refrain from using the ordinary notation U_ℓ as it could cause confusion with the generators of (2.1)). They start as $P_0 = 1$, $P_1 = \beta$ etc. The recursion formula they satisfy, namely $\beta P_\ell = P_{\ell+1} + P_{\ell-1}$, may be used to extend them to $P_{-1} = 0$, $P_{-2} = -1$. We consider these polynomials for θ given by $q = e^{i\theta}$, hence $\beta = q + q^{-1}$.

With these conventions we claim that $\forall L > 0$

$$\sum_{\ell=0}^{\inf(k, L-1)} (-q)^\ell \tilde{Z}_{L-\ell} G_\ell = (-1)^{L-1} P_{L-1}(\beta) G_L \quad (2.21a)$$

$$\sum_{\ell=0}^{\inf(k, L-1)} P_{\ell-2}(\beta) Z_{L-\ell} G_\ell = -P_{L-1}(\beta) G_L. \quad (2.21b)$$

Similar recursion formulae for the physical transfer matrix have been obtained in [15]. Eq. (2.21a) or (2.21b) define recursively all $\tilde{Z}_L$, resp Z_L . With our convention that $G_\ell = 0$ for $\ell > k$, the right hand side of both eqs (2.21) vanishes for $L > k$. Thus (2.21) reads more explicitly

$$L \geq k+1 \quad \tilde{Z}_L = - \sum_{\ell=1}^k (-q)^\ell \tilde{Z}_{L-\ell} G_\ell \quad (2.22a)$$

$$L \leq k \quad \tilde{Z}_L = - \sum_{\ell=1}^{L-1} (-q)^\ell \tilde{Z}_{L-\ell} G_\ell + (-1)^{L-1} P_{L-1}(\beta) G_L \quad (2.22b)$$

and likewise

$$L \geq k+1 \quad Z_L = \sum_{\ell=2}^k P_{\ell-2}(\beta) Z_{L-\ell} G_\ell \quad (2.23a)$$

$$L \leq k \quad Z_L = \sum_{\ell=2}^{L-1} P_{\ell-2}(\beta) Z_{L-\ell} G_\ell + P_{L-1}(\beta) G_L. \quad (2.23b)$$

Note that because of the vanishing of P_1 , the dependence of Z_L on G_1 comes only through that of $Z_1, \dots, Z_k$, and Z_L is thus at most of degree one in G_1 .

The first Z and $\tilde{Z}$ read

$$\begin{aligned}
Z_1 &= G_1 \\
Z_2 &= \beta G_2 \\
Z_3 &= G_1 G_2 + (\beta^2 - 1) G_3 \\
Z_4 &= \beta (G_2^2 + G_1 G_3) + \beta(\beta^2 - 2) G_4 \\
Z_5 &= G_1 G_2^2 + (2\beta^2 - 1) G_2 G_3 + (\beta^2 - 1) G_1 G_4 + (\beta^4 - 3\beta^2 + 1) G_5 \\
&\vdots \qquad \qquad \vdots
\end{aligned} \tag{2.24}$$

$$\begin{aligned}
\tilde{Z}_1 &= G_1 \\
\tilde{Z}_2 &= q G_1^2 - (q + q^{-1}) G_2 \\
\tilde{Z}_3 &= q^2 G_1^3 - (2q^2 + 1) G_1 G_2 + (q^2 + 1 + q^{-2}) G_3 \\
\tilde{Z}_4 &= q^3 G_1^4 - (3q^3 + q) G_1^2 G_2 + (q^3 + q) G_2^2 + (2q^3 + q + q^{-1}) G_1 G_3 \\
&\quad - (q^3 + q + q^{-1} + q^{-3}) G_4 \\
&\vdots \qquad \qquad \vdots
\end{aligned} \tag{2.25}$$

and the reader recognizes in the coefficient of G_L in Z_L resp. $\tilde{Z}_L$ two alternative expressions of $P_{L-1}(\beta)$ (up to a sign).

c) We can also rewrite the right hand sides of (2.24) as *linear* expressions in the N 's, thus getting explicit expressions for (2.16). Curiously, the first polynomials in β turn out to have only positive coefficients. For example the first ones read

$$\begin{aligned}
Z_1 &= N_{\square} \\
Z_2 &= \beta N_{\text{B}} \\
Z_3 &= \beta^2 N_{\text{B}} + N_{\text{BP}} \\
Z_4 &= \beta^3 N_{\text{B}} + \beta (N_{\text{BP}} + N_{\text{B}}) \\
Z_5 &= \beta^4 N_{\text{B}} + \beta^2 (3N_{\text{BP}} + 2N_{\text{B}}) + (N_{\text{BP}} + N_{\text{BP}} + N_{\text{BP}}) \\
&\vdots \qquad \vdots
\end{aligned} \tag{2.26}$$

We have checked that up to Z_9 . The reason of this intriguing feature is not known to us, but should have to do with positivity properties of the operator $U_1 \cdots U_{L-1}$. It also suggests

that the coefficients $z_L^{(\lambda)}$ might be given a combinatorial or group theoretic interpretation.²

Even simpler are the corresponding expressions for the $\tilde{Z}_L$

$$\begin{aligned}
\tilde{Z}_1 &= N_{\square} \\
\tilde{Z}_2 &= qN_{\begin{smallmatrix} \square & \square \end{smallmatrix}} - q^{-1}N_{\begin{smallmatrix} \square \\ \square \end{smallmatrix}} \\
\tilde{Z}_3 &= q^2N_{\begin{smallmatrix} \square & \square & \square \end{smallmatrix}} - N_{\begin{smallmatrix} \square & \square \\ \square \end{smallmatrix}} + q^{-2}N_{\begin{smallmatrix} \square \\ \square & \square \end{smallmatrix}} \\
\tilde{Z}_4 &= q^3N_{\begin{smallmatrix} \square & \square & \square & \square \end{smallmatrix}} - qN_{\begin{smallmatrix} \square & \square & \square \\ \square \end{smallmatrix}} + q^{-1}N_{\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}} - q^{-3}N_{\begin{smallmatrix} \square \\ \square & \square & \square \end{smallmatrix}} \\
\tilde{Z}_5 &= q^4N_{\begin{smallmatrix} \square & \square & \square & \square & \square \end{smallmatrix}} - q^2N_{\begin{smallmatrix} \square & \square & \square & \square \\ \square \end{smallmatrix}} + N_{\begin{smallmatrix} \square & \square & \square \\ \square & \square \end{smallmatrix}} - q^{-2}N_{\begin{smallmatrix} \square & \square \\ \square & \square & \square \end{smallmatrix}} + q^{-4}N_{\begin{smallmatrix} \square \\ \square & \square & \square & \square \end{smallmatrix}}
\end{aligned} \tag{2.27}$$

These formulae suggest an obvious generalization. Consider the Young tableaux with at most one line of length larger than one (i.e. of type $(t+1)(1)^s$, $s+t+1=L$, $s, t \geq 0$) and denote by $\lambda_{t,s}$ the corresponding weight of $sl(k)$ ($k \geq s+1$)

$$\lambda = \lambda_{t,s} := \begin{array}{c} \overbrace{\square \square \square \square \square \square \square \square}^{t+1} \\ \square \\ \square \\ \square \end{array} \Bigg\}^s$$

Then

$$\tilde{Z}_L = \sum_{\substack{\lambda_{t,s} \\ s+t+1=L}} (-1)^s q^{L-1-2s} N_{\lambda_{t,s}} \tag{2.28}$$

As we shall see in sect. 3.1, this conjectured expression is indeed the solution to eq. (2.21a) or (2.22) and it reduces to a well known fact in the classical limit $q \rightarrow 1$.

d) Explicit solutions to eqs (2.21b) may be given for $k=2, 3$. For $k=2$,

$$\begin{aligned}
Z_{2\ell+1} &= G_1 G_2^\ell \\
Z_{2\ell} &= \beta G_2^\ell
\end{aligned} \tag{2.29}$$

² Here again, to get universal expressions, we have extended slightly the original conventions on the weights or Young tableaux labelling the fusion matrices N : for a given k , any column in a Young tableau with k rows may be deleted, any tableau with a column of more than k rows does not contribute. Thus for instance $N_{\begin{smallmatrix} \square \\ \square \end{smallmatrix}} = \mathbb{I}$ if $k=4$, and $=0$ for $k \leq 3$.

For $k = 3$,

$$\begin{aligned}
Z_{2\ell+1} &= \sum_{p=0}^{\lfloor \frac{\ell}{3} \rfloor} \binom{\ell-p}{2p} \beta^{2p} G_1 G_2^{\ell-3p} G_3^{2p} \\
&\quad + \sum_{p=0}^{\lfloor \frac{\ell-1}{3} \rfloor} \left[\binom{\ell-p}{2p+1} \beta^{2p+2} - \binom{\ell-p-1}{2p} \beta^{2p} \right] G_2^{\ell-3p-1} G_3^{2p+1} \\
Z_{2\ell} &= \beta G_2^\ell + \sum_{p=1}^{\lfloor \frac{\ell}{3} \rfloor} \left[\binom{\ell-p}{2p} \beta^{2p+1} - \binom{\ell-p-1}{2p-1} \beta^{2p-1} \right] G_2^{\ell-3p} G_3^{2p} \\
&\quad + \sum_{p=0}^{\lfloor \frac{\ell-2}{3} \rfloor} \binom{\ell-p-1}{2p+1} \beta^{2p+1} G_1 G_2^{\ell-3p-2} G_3^{2p+1}
\end{aligned} \tag{2.30}$$

In these two cases $k = 2, 3$, we have respectively $G_2 \equiv 1$, $G_3 \equiv 1$, but we have left them in order to make explicit the homogeneity of Z_ℓ . In checking that these expressions do satisfy the recursion formulae (2.23) it is useful to use the following formula for $P_n(\beta)$:

$$P_n(\beta) = \sum_{p=0}^{\lfloor \frac{n}{2} \rfloor} (-1)^p \binom{n-p}{p} \beta^{n-2p} . \tag{2.31}$$

Similar expressions may be obtained for higher k but they become increasingly cumbersome. Eq. (2.29) may be easily proved in full generality for all graphs using Pasquier's expressions for the Boltzmann weights [2].

e) The status of the assertions (2.21) is as follows

- 1) the expression (2.28) may be proved to solve (2.21a), and its limit as $q \rightarrow 1$ is related to well known properties of the symmetric group; as a result, (2.21a) itself is proved in that limit $q = 1$ (see below, sect. 3.1);
- 2) eq. (2.21a) may be shown to be a simple consequence of (2.3) for $k = 2, 3$ and all $L > k$ (see below, sect. 3.2); thus it holds for *all* graphs;
- 3) eq. (2.21b) has been checked for the basic graph of $sl(k)$ up to $k = 9$ and $L = 9$ using the explicit expression (2.10);
- 4) eq. (2.21b) has been proved for the basic graph of $sl(k)$, $k = 3, 4$, all $L > 0$ (see below, sect. 3.3);
- 5) the non trivial equivalence between (2.21a) and (2.21b) has been verified for $k = 2, 3, 4$;

6) these identities are consistent with another identity, related to the Markov property of the trace (see footnote 1), namely that

$$\sum_c \begin{array}{c} c \\ \diagup \quad \diagdown \\ b \quad \quad b \\ \diagdown \quad \diagup \\ a \end{array} \psi_c^{(1)} = P_{k-2}(\beta) \psi_b^{(1)} , \quad (2.32)$$

which may be proved on the expression (2.10). (For other graphs, it then follows from the existence of ‘cells’ [16,4,17,18]). From that identity, it follows that

$$\langle Z_L \rangle := \sum_{a,b} \psi_a^{(1)} (Z_L)_{ab} \psi_b^{(1)} = P_{k-1}(\beta) P_{k-2}^{L-1}(\beta) , \quad (2.33)$$

and it is easy to see that (2.33) satisfies the recursion formula (2.21b).

In view of these evidences, it seems quite reasonable to conjecture the general validity of (2.21) for all representations, all k and all $L > k$.

3. Elements of proof.

In this section we collect the various evidences on the validity of eqs (2.21) that we have found

3.1. (2.28) solves (2.22); the $q \rightarrow 1$ limit of eq. (2.22)

First let us consider the expression (2.28) and let us show why it is natural in the limit where $n \rightarrow \infty$, hence $q \rightarrow 1$. Then the paths we consider in the basic case are just those on the entire weight lattice of $sl(k)$. For q , the deformation parameter equal to 1, the Hecke algebra $\mathcal{H}_L$ reduces to the algebra of the symmetric group S_L and its irreducible representations associated with paths of length L from $\mathbf{1}$ to λ are nothing else than the representations of S_L attached to the Young tableau associated with λ . Thus for any $\sigma \in \mathcal{H}_L$, the matrix element $(\text{tr } \sigma)_{\mathbf{1}\lambda}$ is simply the irreducible character $\chi_\lambda(\sigma)$ of S_L . Moreover, the product $g_1 g_2 \dots g_{L-1}$ whose trace defines $\tilde{Z}_L$ in (2.13) is a cyclic permutation of S_L . It is well known (see for example [19]) that the only Young tableaux giving non vanishing characters of cyclic permutations σ of S_L are those denoted $\lambda_{t,s}$ in sect. 2.4.c, for which $\chi_{\lambda_{t,s}}(\sigma) = (-1)^s$. Thus in the limit $q = 1$,

$$(\tilde{Z}_L)_{\mathbf{1}\lambda} = \sum_{\lambda_{t,s}} (-1)^s \delta_{\lambda \lambda_{st}} \quad (3.1)$$

or


$$\tilde{Z}_L = \sum_{\substack{\lambda_{t,s} \\ s+t+1=L}} (-1)^s N_{\lambda_{t,s}} , \quad (2.28)_1$$

and expression (2.28) is a very natural q -deformation of that formula.

In fact one may prove that (2.28) satisfies (2.22) for all q . Considering first the case where $k \geq L$, insert (2.28) in the $(1, \mu)$ matrix element of the l.h.s. of (2.21a).

$$\sum_{\lambda} \sum_{\ell=0}^{L-1} (-q)^\ell (\tilde{Z}_{L-\ell})_{1\lambda} (G_\ell)_{\lambda\mu} \stackrel{?}{=} \sum_{\ell=0}^{L-1} \sum_{\substack{\lambda_{t,s} \\ s+t+1=L-\ell}} (-1)^s q^{L-1-\ell-2s} (-q)^\ell (G_\ell)_{\lambda_{t,s}\mu} \quad (3.2)$$

It receives contributions from those weights μ that appear in the tensor product of $\lambda_{t,s}$ by the ℓ -th fundamental representation



It is then a simple exercise of bookkeeping to see that there are cancellations for almost all μ and that only the vertical Young tableau with L boxes contributes $(-1)^{L-1}(q^{L-1} + q^{L-3} + \dots + q^{-(L-1)}) = (-1)^{L-1}P_{L-1}(\beta)$ in agreement with the conjectured formula. For example, for $L = 3$, $k \geq 3$

$$\begin{aligned} \tilde{Z}_3 N_0 - q \tilde{Z}_2 N_1 + q^2 \tilde{Z}_1 N_2 &= (q^2 \square\square\square - \boxplus + q^{-2} \boxminus) - q (q\square\square - q^{-1} \boxplus) \otimes \square + q^2 \square \otimes \boxplus \\ &= q^2 \square\square\square - \boxplus + q^{-2} \boxminus - q^2 \square\square - q^2 \boxplus + \boxplus + \boxminus + q^2 \boxplus + q^2 \boxminus \quad (3.3) \\ &= (q^2 + 1 + q^{-2}) \boxminus . \end{aligned}$$

For $k < L$, the same discussion applies, provided one restricts oneself to Young tableaux with less than $k + 1$ rows. For instance, for $k = 2$, the contribution of $\boxminus$ in (3.3) has to be removed.

In this section, we have proved that i) eq. (2.28) is true for $q = 1$; ii) that it satisfies (2.21a) for all q . This implies that the conjectured identity (2.21a) is itself established for $q = 1$.

3.2. (2.3) implies (2.21a)

This implication has been checked for $k = 2, 3$. For $k = 3$, to write explicitly (2.3), we have to enumerate all permutations of the symmetric group S_4 and to decompose them

into products of adjacent transpositions g_i , $i = 1, 2, 3$. We get in that way

$$\begin{aligned}
& I - q^{-1}(g_1 + g_2 + g_3) + q^{-2}(g_1g_3 + g_2g_1 + g_1g_2 + g_2g_3 + g_3g_2) \\
& - q^{-3}(g_1g_2g_1 + g_2g_3g_2 + g_3g_2g_1 + g_1g_3g_2 + g_1g_2g_3 + g_2g_3g_1) \\
& + q^{-4}(g_2g_1g_3g_2 + g_3g_1g_2g_3 + g_3g_2g_1g_3 + g_1g_2g_3g_1 + g_1g_3g_2g_1) \\
& - q^{-5}(g_2g_3g_2g_1g_2 + g_2g_3g_1g_2g_3 + g_1g_2g_3g_2g_1) + q^{-6}g_2g_1g_2g_3g_2g_1 = 0 .
\end{aligned} \tag{3.4}$$

Multiply this by $g_4g_5 \cdots g_{L-1}$, take the trace and use eqs (2.1) and the cyclicity of the trace to simplify this expression. A straightforward but tedious calculation yields

$$(1 + q^{-2})(q^{-3} + q^{-5} + q^{-7}) \left(-\tilde{Z}_L + q\tilde{Z}_{L-1}G_1 - q^2\tilde{Z}_{L-2}G_2 + q^3\tilde{Z}_{L-3}G_3 \right) = 0 \tag{3.5}$$

which is the appropriate eq. (2.21a). The same can be done for $k = 2$ and might be repeated for all $k > 3$, although we have not been able to master the combinatorics.

3.3. Proving (2.21b) for all L and $k = 3, 4$

For the basic graph for $k = 3, 4$, using the explicit expressions (2.10) of the matrix elements of the U 's, we have been able to prove (2.21b) for all $L \geq k$ (while it is verified for $L \leq k$ by simple inspection). The proof displays an interesting and suggestive hierarchical structure.

Let us sketch the proof for $k = 3$. The identity to be proved reads

$$Z_L = Z_{L-2} + \beta Z_{L-3} . \tag{3.6}$$

Denote $a \xrightarrow{\alpha} b$ an edge between the two heights a and b of the “basic graph” (a and b are integrable weights) along the direction e_α . We look at the contributions to Z_L from paths ending at some height e that passed through the heights a and b , i.e. paths

$$a \xrightarrow{\alpha} b \xrightarrow{\beta} c \xrightarrow{\delta} d \xrightarrow{\epsilon} e . \tag{3.7}$$

with a, b, α, e fixed, and we shall sum over the possible c, d . The direction β must be different from α and *by convention*, γ denotes the third direction such that $e_\alpha + e_\beta + e_\gamma = 0$.

Two cases are possible:

- 1) β, δ and ϵ are all different, hence equal up to some permutation to α, β, γ , hence $e = b$.

This is possible for

$$a \xrightarrow{\alpha} b \xrightarrow{\beta} c \xrightarrow{\alpha} d \xrightarrow{\gamma} e = b$$

or for

$$a \xrightarrow{\alpha} b \xrightarrow{\beta} c \xrightarrow{\gamma} d \xrightarrow{\alpha} e = b ;$$

2) β, δ, ϵ not all different, either as

$$a \xrightarrow{\alpha} b \xrightarrow{\beta} c \xrightarrow{\gamma} d \xrightarrow{\beta} e = c + e_{\beta} + e_{\gamma}$$

or as

$$a \xrightarrow{\alpha} b \xrightarrow{\beta} c \xrightarrow{\alpha} d \xrightarrow{\beta} e = c + e_{\beta} + e_{\alpha} .$$

Let us now compute the contributions of these paths to the l.h.s. of the recursion formula

$$\begin{aligned} \sum_{c,d} Z_{ab}^{(L-3)} & \begin{array}{c} c \\ \diamond \\ a \end{array} b \begin{array}{c} d \\ \diamond \\ b \end{array} c \begin{array}{c} e \\ \diamond \\ c \end{array} d = \sum_j Z_{ab}^{(L-3)} \begin{array}{c} b+e_{\beta} \\ \diamond \\ a \end{array} b \\ & \left[\begin{array}{c} d=c+e_{\gamma} \\ c=b+e_{\beta} \diamond c \end{array} d \begin{array}{c} b \\ \diamond \\ c \end{array} d + \begin{array}{c} d=c+e_{\alpha} \\ c=b+e_{\beta} \diamond c \end{array} d \begin{array}{c} b \\ \diamond \\ c \end{array} d \right. \\ & \left. + \begin{array}{c} d=c+e_{\gamma} \\ c=b+e_{\beta} \diamond b \end{array} c \begin{array}{c} e=c+e_{\beta}+e_{\gamma} \\ \diamond \\ c \end{array} d + \begin{array}{c} d=c+e_{\alpha} \\ c=b+e_{\beta} \diamond b \end{array} c \begin{array}{c} e=c+e_{\beta}+e_{\alpha} \\ \diamond \\ c \end{array} c \right] \end{aligned} \quad (3.8)$$

where $Z_{ab}^{(L-3)}$ is the contribution to Z_{L-2} from the paths ending along the edge ab .

The last two terms in the square bracket are equal to one. Indeed with the notations of (2.10)

$$\begin{array}{c} d=c+e_{\gamma} \\ c=b+e_{\beta} \diamond b \end{array} c \begin{array}{c} e=c+e_{\beta}+e_{\gamma} \\ \diamond \\ c \end{array} d = \frac{s_{\beta\gamma}(c)}{s_{\beta\gamma}(b)} \frac{s_{\gamma\beta}(d)}{s_{\gamma\beta}(c)} = 1 \quad (3.9)$$

since $c = b + e_{\beta}$ hence $s_{\gamma\beta}(d) = s((e_{\gamma} - e_{\beta}), (b + e_{\gamma} + e_{\beta})) = -s_{\beta\gamma}(b)$ and $s_{\beta\gamma}(c) = -s_{\gamma\beta}(c)$.

Let us compare the result of (3.8) with the contributions to the r.h.s. of (3.7) of the same paths through a and b

$$Z_{ab}^{(L-2)} \left[\sum_j \begin{array}{c} c=b+e_{\beta} \\ b \diamond b \\ a \end{array} (G_2)_{ce} + \beta \delta_{be} \right] . \quad (3.10)$$

Since $(G_2)_{ce} = \delta_{e,c+e_\alpha+e_\beta} + \delta_{e,c+e_\beta+e_\gamma} + \delta_{e,c+e_\gamma+e_\alpha}$, the first two terms reproduce the last two in (2), and the last is equal to δ_{eb} , hence we have to compare all the terms in (3.8) and (3.10) proportional to δ_{be} . Let us show indeed that (i being fixed, *not* summed over)

$$\sum_{\beta \neq \alpha} b \begin{array}{c} c=b+e_\beta \\ \diamond \\ a \end{array} b=a+e_\alpha \left[c=b+e_\beta \begin{array}{c} d=c+e_\gamma \\ \diamond \\ b \end{array} c \begin{array}{c} b \\ \diamond \\ c \end{array} d + c=b+e_\beta \begin{array}{c} d=c+e_\alpha \\ \diamond \\ b \end{array} c \begin{array}{c} b \\ \diamond \\ c \end{array} d - 1 \right] = \beta. \quad (3.11)$$

The l.h.s. of (4) reads, if $a_{\alpha\beta} := (a, e_\alpha - e_\beta)$ and $s(x) = \sin \frac{\pi x}{n}$

$$\begin{aligned} l.h.s. &= \sum_{j \neq i} \frac{s(a_{\alpha\beta} + 1)}{s(a_{\alpha\beta})} \left[\frac{s(b_{\beta\gamma} + 1)}{s(b_{\beta\gamma})} \frac{s(c_{\gamma\alpha} + 1)}{s(c_{\gamma\alpha})} + \frac{s(b_{\beta\alpha} + 1)}{s(b_{\beta\alpha})} \frac{s(c_{\alpha\gamma} + 1)}{s(c_{\alpha\gamma})} - 1 \right] \\ &= \sum_{j \neq i} \left[\frac{s(a_{\alpha\beta} + 1)}{s(a_{\alpha\beta})} \frac{s(a_{\beta\gamma} + 1)}{s(a_{\beta\gamma})} \frac{s(a_{\gamma\alpha})}{s(a_{\gamma\alpha} - 1)} + \frac{s(a_{\alpha\gamma} + 2)}{s(a_{\alpha\gamma} + 1)} - \frac{s(a_{\alpha\beta} + 1)}{s(a_{\alpha\beta})} \right] \\ &= \frac{s(a_{\alpha\beta} + 1)}{s(a_{\alpha\beta})} \frac{s(a_{\beta\gamma} + 1)}{s(a_{\beta\gamma})} \frac{s(a_{\gamma\alpha})}{s(a_{\gamma\alpha} - 1)} + \frac{s(a_{\alpha\gamma} + 2)}{s(a_{\alpha\gamma} + 1)} - \frac{s(a_{\alpha\beta} + 1)}{s(a_{\alpha\beta})} \\ &\quad + \frac{s(a_{\alpha\gamma} + 1)}{s(a_{\alpha\gamma})} \frac{s(a_{\gamma\beta} + 1)}{s(a_{\gamma\beta})} \frac{s(a_{\beta\alpha})}{s(a_{\beta\alpha} - 1)} + \frac{s(a_{\alpha\beta} + 2)}{s(a_{\alpha\beta} + 1)} - \frac{s(a_{\alpha\gamma} + 1)}{s(a_{\alpha\gamma})}. \end{aligned} \quad (3.12)$$

One may check by a direct (and boring) calculation that this is equal to β but it is simpler to regard it as a function of the components a_1 and a_2 of a ; this is a meromorphic function; it is bounded as a_1, a_2 go to ∞ , and the residues at the possible poles ($a_{\alpha\beta} = 0 \bmod n$, $a_{\beta\gamma} = 0 \bmod n$, $a_{\alpha\gamma} = -1 \bmod n$) all vanish. By Liouville theorem, we conclude that this is a constant, that may be computed by letting a_1, a_2 go to $-i\infty$, for which we find the result β , which establishes (3.11) and hence (3.7).

We have repeated the same analysis for the case of $sl(4)$, i.e. $k = 4$. What is quite intriguing is that the identity (3.12) plays a role several times in that $k = 4$ calculation: first to simplify some contributions, in the same way as (3.9) did, and secondly to help to establish the analogue of (3.12). The relevant (very cumbersome) trigonometric identity is again proved by making use of Liouville theorem, and the previous identity (3.12) is just what guarantees the vanishing of the residues at the possible poles. There is little doubt that this structure extends to higher k but we have not been able to fully disentangle the hierarchy of identities that appear there.

Note that this approach hints at a property apparently stronger than the identities (2.21), namely that, for $k = 3$ say,

$$(Z_4 - Z_2 G_2 - \beta Z_1 G_3)_{ab;e} = 0 \quad (3.13)$$

where $(Z_L)_{ab;e}$ denotes a trace like in (2.13) but with two heights fixed on one side, one on the other. Such partition functions have been already the object of some interest in the past [8,9].

4. Conclusions and perspectives

In this paper we have presented some evidence that traces of operators in representations of the Hecke algebra satisfy remarkable identities that have a large character of universality, that is, that are independent of the representation. Clearly a more systematic proof is still needed. Also one should extend it to cases of graphs with multiple edges (as those appearing in orbifold graphs [3]); we do have evidence that the identities remain valid in those cases, when the appropriate modifications, v.i.z. the introduction of additional *edge* degrees of freedom, are taken into account.

Let us mention some directions in which these identities are or might be useful:

- As mentionned in the Introduction, these traces also yield contributions to the two-point function of lattice operators, the so-called “Pasquier operators”: cf [2,10]. It is an interesting question to see if one can extend this to 3-point functions, as it would provide some non trivial information on the operator algebra. This has been achieved so far only for $k = 2$ [10], and there is some indication that the situation for $k \geq 3$ is more subtle.

- Regarding the sequences of U or g appearing in the traces (2.13) as the product along a cylinder, one can also use the exact results of this paper to compute the partition function of a quasi-one-dimensional system. We refer the interested reader to [15] for a detailed discussion.

- The original hope was that the detailed knowledge of these identities would provide additional and sufficient information to fully determine the Boltzmann weights in cases where they are not known. This is based on the following observation.

A powerful method of determination of Boltzmann weights is through the calculation of “cells” [16,4,17,18], that intertwine between the weights pertaining to the graph $\mathcal{G}_1$ and those relative to some basic graph $\mathcal{A}$. Some identities, like (2.32), are such that if they are satisfied by the basic graphs, the mere existence of cells (with their orthonormality properties) guarantees that they are also satisfied by the graph $\mathcal{G}_1$. This is *not* the case with identities (2.21). Thus if these identities are satisfied both by $\mathcal{A}$ and $\mathcal{G}_1$, this implies non trivial constraints on the cells. At this stage, the systematic exploitation of these constraints has not yet been achieved. We hope to report on that matter in the future.

Acknowledgements It is a pleasure to acknowledge stimulating discussions with P. Di Francesco, V. Pasquier and P. Pearce. Y.-K. Z. is supported by the Australian Research Council and partially by Natural Science Foundation of China. J.-B. Z. wants to thank

O. Foda and the Mathematics Department, University of Melbourne, for their generous hospitality that made this collaboration possible.

Appendix A.

This appendix presents the proof of the assertion of eq. (2.16), in the case of the “basic” graphs $\mathcal{A}^{(n)}$.

Let $V^{(\lambda)}$ denote the representation space of weight λ of the algebra $\mathcal{U}_q sl(k)$. We recall that for q is a root of unity, $q = \exp \frac{i\pi}{n}$, it is consistent to restrict to the admissible weights $\lambda \in \mathcal{P}_{++}^{(n)}$. If W is a certain representation space of the quantum group, we thus decompose W as $W \cong \oplus_{\lambda} \mathcal{C}_W^{(\lambda)} \otimes V^{(\lambda)}$, with multiplicity spaces $\mathcal{C}_W^{(\lambda)}$, and denote $\mathbf{1}_W^{(\lambda)}$ the projector on the irreducible representation (λ) . By Schur’s lemma any operator x on W that commutes with the action of the quantum group reads $x = \oplus_{\lambda} x^{(\lambda)} \otimes \mathbf{1}_W^{(\lambda)}$.

To describe the subspace $\mathcal{P}_{\lambda_0 \lambda_L}^{(L)}$ of paths of length L on a basic graph going from λ_0 to λ_L , it is appropriate to proceed as follows

- i) one introduces the space $V_0 = V^{(\lambda_0)}$ and attach it to site 0 of the chain,
- ii) one introduces L copies $V_1, \dots, V_L$ of the k -dimensional representation of $\mathcal{U}_q sl(k)$, attached to the successive *edges* of the path.

If one decomposes

$$V_0 \otimes V_1 \otimes \dots \otimes V_L = \oplus_{\lambda} \mathcal{C}^{(\lambda)} \otimes V^{(\lambda)} , \quad (\text{A.1})$$

then

$$\mathcal{P}_{\lambda_0 \lambda_L}^{(L)} = \mathcal{C}^{(\lambda_L)} . \quad (\text{A.2})$$

Implicit in this procedure is the connection between height (IRF) models and “vertex models” where the spins take their values in some fixed representation of $\mathcal{U}_q sl(k)$ (here the fundamental, k -dimensional one) [20].

Let us denote $W := V_1 \otimes \dots \otimes V_L$ for short and let $\Delta(\mathbf{1}^{(\lambda_L)})$ be the projection on the irreducible representation (λ_L) induced by the coproduct in the space $V_0 \otimes W$. We are interested in operators x that are generated by the Hecke algebra and thus are in the commutant of the quantum group acting on W . Also these x act as the identity on V_0 . For such an operator, computing the trace in the space $V_0 \otimes W$ of $(\mathbf{1}_0^{(\lambda_0)} .x. \Delta(\mathbf{1}^{(\lambda_L)}))$ amounts to taking the trace in the subspace $\mathcal{P}_{\lambda_0 \lambda_L}^{(L)}$

$$\text{tr}_{V_0 \otimes \dots \otimes V_L} (\mathbf{1}_{V_0}^{(\lambda_0)} .x. \Delta(\mathbf{1}^{(\lambda_L)})) = \text{tr}_{\mathcal{P}_{\lambda_0 \lambda_L}^{(L)}} (x) \text{tr} (\mathbf{1}^{(\lambda_L)}) , \quad (\text{A.3})$$

and $\text{tr}_{\mathcal{P}_{\lambda_0 \lambda_L}^{(\lambda)}}(x)$ is what is denoted $(\text{tr}(x))_{\lambda_0 \lambda_L}$ in eq. (2.12); $\text{tr} \mathbf{1}^{(\lambda_L)}$ is the dimension of the irreducible representation (λ_L) .

By the remarks above, one may decompose $W = \oplus_{\lambda} \mathcal{C}_W^{(\lambda)} \otimes V^{(\lambda)}$, write accordingly $x = \text{id}_0 \otimes \sum_{\lambda} (x^{(\lambda)} \otimes \mathbf{1}^{(\lambda)})$, and compute

$$\begin{aligned} (\text{tr}(x))_{\lambda_0 \lambda_L} \text{tr}(\mathbf{1}^{(\lambda_L)}) &= \sum_{\lambda \in \mathcal{P}_{++}^{(n)}} \text{tr}_{V_0 \otimes (\mathcal{C}_W^{(\lambda)} \otimes V^{(\lambda)})} \left(\mathbf{1}_0^{(\lambda_0)} \cdot (x^{(\lambda)} \otimes \mathbf{1}_W^{(\lambda)}) \cdot \Delta(\mathbf{1}^{(\lambda_L)}) \right) \\ &= \sum_{\lambda \in \mathcal{P}_{++}^{(n)}} \text{tr}_{V_0 \otimes W} \left(\mathbf{1}_{V_0}^{(\lambda_0)} \cdot \mathbf{1}_W^{(\lambda)} \cdot \Delta(\mathbf{1}^{(\lambda_L)}) \right) \text{tr}_{\mathcal{C}_W^{(\lambda)}} x^{(\lambda)}. \end{aligned} \quad (\text{A.4})$$

The first trace is nothing else than the fusion coefficient $N_{\lambda_0 \lambda}^{\lambda_L}$ times $\text{tr}(\mathbf{1}^{(\lambda_L)})$. If we call $x_L^{(\lambda)}$ the last trace appearing in (A.4), we obtain the desired result (2.16) for the basic graph, for which $N_{\lambda}^{(\mathcal{A})} \equiv N_{\lambda}$.

References

- [1] N.Yu. Reshetikhin, *Quantized Universal Enveloping Algebras, the Yang-Baxter Equation and Invariants of Links*, I and II, LOMI preprints 1988.
- [2] V. Pasquier, *Nucl. Phys.* **B285** [FS19] (1987) 162 ;
V. Pasquier, *J. Phys.* **A20** (1987) 5707.
- [3] I.K. Kostov, *Nucl. Phys.* **B 300** [FS22] (1988) 559.
- [4] P. Di Francesco and J.-B. Zuber, *Nucl. Phys.* **B338** (1990) 602.
- [5] P. Di Francesco and J.-B. Zuber, in *Recent Developments in Conformal Field Theories*, Trieste Conference 1989, S. Randjbar-Daemi, E. Sezgin and J.-B. Zuber eds., World Scientific 1990 ;
P. Di Francesco, *Int. J. Mod. Phys.* **A7** (1992) 407.
- [6] N. Sochen, *Nucl. Phys.* **B360** (1991) 613.
- [7] V.B. Petkova and J.-B. Zuber, [hep-th 9510175](#), *Nucl. Phys. B* **B463** (1996) 161.
- [8] H. Saleur and M. Bauer, *Nucl. Phys.* **B320** (1989) 591.
- [9] J. Cardy, *Nucl. Phys.* **B324** (1989) 581.
- [10] V.B. Petkova and J.-B. Zuber, *Nucl. Phys.* **B438** (1995) 347.
- [11] V.V. Bazhanov and N.Yu. Reshetikhin, *J. Phys.* **A 23** (1990) 1477.
- [12] E. Verlinde, *Nucl. Phys.* **B300** [FS22] (1988) 360.
- [13] H. Wenzl, *Representations of Hecke Algebras and Subfactors*, Ph.D. Thesis, University of Pennsylvania.
- [14] V. Pasquier and H. Saleur, *Nucl. Phys.* **B330** (1990) 523.
- [15] S. Loesch, *The exact solution of quasi one-dimensional Hecke algebra models*, Diplomarbeit in Physik, Universität Bonn report, November 1995.
- [16] A. Ocneanu, *Quantized groups, string algebras and Galois theory for algebras in Operator Algebras and Applications*, D.E. Evans and M. Takesaki eds, Lond. Math. Soc. Lecture Note Series **136**, Cambridge University Press, 119.
- [17] P. Roche, *Comm. Math. Phys.* **127** (1990) 395.
- [18] P. Pearce and Y.-K. Zhou, *Int. J. Mod. Phys. B* **7** (1993) 3649.
- [19] C. Itzykson and J.-B. Zuber, *Int. J. Mod. Phys.* **A7** (1992) 5661 and earlier references therein.
- [20] V. Pasquier, CEnology of IRF models, *Comm. Math. Phys.* **118** (1988) 355.